Lactate dehydrogenase isoenzymes in dimethylnitrosamine-induced hepatic fibrosis in rats

By: George, J (George, J); Chandrakasan, G (Chandrakasan, G)

View ResearcherID and ORCID

JOURNAL OF CLINICAL BIOCHEMISTRY AND NUTRITION

Volume: 22 Issue: 1 Pages: 51-62

Published: JAN 1997 **Document Type: Article** View Journal Impact

Abstract

Alteration of liver function is an inevitable phenomenon in hepatic fibrosis. Since many enzymes are useful in diagnosing liver diseases, the alteration of the lactate dehydrogenase (LDH) isoenzyme pattern was studied in experimentally induced liver fibrosis. The liver injury was induced by intraperitoneal injections of dimethylnitrosamine (DMN) into adult male albino rats belonging to the Wistar strain. The injections were given on the first 3 consecutive days of each week over a period of 3 weeks. The degree of hepatic fibrosis was assessed by histopathological examination as well as by monitoring the collagen content of the liver tissue. The LDH isoenzymes in serum and liver were separated by polyacrylamide gel electrophoresis. The percentage distribution of the isoenzymes was determined, and the M/H ratio was calculated. The total LDH activity was also studied in the liver and serum samples. The results demonstrated many pathological changes including centrilobular necrosis and fibrosis after DMN treatment. The collagen content was increased to about 4 times normal by the 21st day of DMN administration. The total LDH activity was significantly increased in the serum on the 14th and 21st days. But in the liver tissue, a significant increase was noticed only on the 7th day. The LDH isoenzyme patterns indicated a sharp increase in the activities of LDH, and LDH, in the serum on the 14th and 21st days of DMN treatment. A remarkable increase was noticed in the M/H ratio on the 14th and 21st days in the serum and on all the days in the liver tissue. Increased serum LDH activity suggests deterioration of liver functions during hepatic fibrosis. The present study demonstrated that serum LDH isoenzyme assay is a useful tool in the diagnosis of hepatic fibrosis along with other biochemical tests.

Keywords

Author Keywords: lactate dehydrogenase; LDH isoenzymes; dimethylnitrosamine; hepatic fibrosis; liver collagen

KeyWords Plus: INDUCED CIRRHOSIS; LIVER-DISEASE; INJURY; SERUM; CELLS

Author Information

Addresses:

+ [1] CENT LEATHER RES INST, DEPT BIOCHEM, MADRAS 600020, TAMIL NADU, INDIA

Publisher

JOURNAL CLINICAL BIOCHEMISTRY & NUTRITION, KYOTO PREFECTURAL UNIV MED, GRAD SCH MEDICAL SCIENCE, DEPT MOLECULAR GASTROENTEROLOGY & HEPATOLOGY, KYOTO, 602-8566, JAPAN

Journal Information

Impact Factor: Journal Citation Reports

Categories / Classification

Research Areas: Nutrition & Dietetics

Web of Science Categories: Nutrition & Dietetics

See more data fields

Citation Network

In Web of Science Core Collection

13

Times Cited



Create Citation Alert

All Times Cited Counts

13 in All Databases

See more counts

31

Cited References

View Related Records

Most recently cited by:

Panchal, Shital S.; Ghatak, Somsuvra B.; Jha, Abhishek B.; et al. Reduction of liver tumerogenic effect of N-nitrosodiethylamine by treatment with gamma-oryzanol in Balb/C mice. **ENVIRONMENTAL TOXICOLOGY AND** PHARMACOLOGY (2017)

Arulkumar, Mani; Vijayan, Raji; Penislusshiyan, Sakayanathan; et al. Alteration of paraoxonase, arylesterase and lactonase activities in people around fluoride endemic area of Tamil Nadu,

CLINICA CHIMICA ACTA (2017)

View All

Use in Web of Science

Web of Science Usage Count

0

Last 180 Days

Since 2013

Learn more

This record is from: Web of Science Core Collection Science Citation Index Expanded