

Elevation of Serum Levels of Advanced Glycation End Products in Patients With Non-B or Non-C Hepatocellular Carcinoma

Hiromi Kan,¹ Sho-ichi Yamagishi,^{2*} Ayako Ojima,² Kei Fukami,³ Seiji Ueda,³

Masayoshi Takeuchi,⁴ Hideyuki Hyogo,¹ Hiroshi Aikata,¹ and Kazuaki Chayama¹

¹Department of Gastroenterology and Metabolism, Hiroshima University Hospital, Hiroshima, Japan

²Department of Pathophysiology and Therapeutics of Diabetic Vascular Complications, Kurume University School of Medicine, Kurume, Japan

³Department of Medicine, Kurume University School of Medicine, Kurume, Japan

⁴Department of Advanced Medicine, Medical Research Institute, Kanazawa Medical University, Kanazawa, Japan

Background: The prevalence of non-B or non-C hepatocellular carcinoma (NBNC-HCC) has been increasing all over the world. Advanced glycation end products (AGE) play a role in the pathogenesis of alcoholic liver injury or nonalcoholic steatohepatitis (NASH). **Methods:** We examined here whether serum levels of AGE were elevated in NBNC-HCC patients compared with NASH subjects without HCC and investigated which anthropometric and clinical variables were independent determinants of AGE. **Results:** Ninety NBNC-HCC, 56 NASH, and 27 control subjects underwent a complete history and physical examination, determination of blood chemistries, including AGE levels. Serum levels of AGE were significantly higher in NBNC-HCC patients compared with NASH and control subjects

[9.1 ± 2.7 , 5.2 ± 1.7 , 3.5 ± 1.2 (U/ml), respectively, $P < 0.05$]. Univariate analysis showed that AGE levels were associated with male ($P < 0.05$), age ($P < 0.01$), aspartate aminotransferase ($P < 0.05$), γ -glutamyl transpeptidase (GGT) ($P < 0.01$), HDL-cholesterol (inversely, $P < 0.01$), fasting plasma glucose ($P < 0.01$), and HbA1c ($P < 0.05$). By the use of multiple stepwise regression analysis, age, GGT, and HDL-cholesterol (inversely) remained significant and were independently related to AGE levels ($R^2 = 0.406$). **Conclusion:** The present results suggest that AGE might be involved in the pathogenesis of NBNC-HCC, thereby being a biomarker that could discriminate NBNC-HCC from NASH. J. Clin. Lab. Anal. 29:480–484, 2015. © 2014 Wiley Periodicals, Inc.

Key words: AGE; NASH; Non-B or non-C hepatocellular carcinoma

INTRODUCTION

Hepatocellular carcinoma (HCC) is one of the most common malignancies and causes of cancer-related deaths in the world (1,2). Although most cases of HCC are attributable to chronic liver disease resulting from chronic hepatitis B virus (HBV) or hepatitis C virus (HCV) infections, a substantial proportion of HCC patients are negative for markers of HBV surface antigen (HBVAg) or HCV antibody (HCVAb), being diagnosed as non-B or non-C HCC (NBNC-HCC). The frequency of NBNC-HCC has been reported to be 5–15%, and number of NBNC-HCC has been increasing gradually all over the world (3–8). Alcoholic or metabolic derangement-related liver damage could contribute to the pathogenesis of NBNC-

HCC (3–8). However, in contrast to HBV- or HCV-related HCC, clinical characteristics of NBNC-HCC are not fully examined. Therefore, carcinomas in these patients were

Grant sponsor: Grants-in-Aid for Scientific Research (B) from the Ministry of Education, Culture, Sports, Science, and Technology, Japan (to M.T. and to S.Y.); Grant sponsor: Strategic Research Foundation at Private Universities, the Ministry of Education, Culture, Sports, Science, and Technology (MEXT, to S.Y.).

*Correspondence to: Sho-ichi Yamagishi, Department of Pathophysiology and Therapeutics of Diabetic Vascular Complications, Kurume University School of Medicine, 67 Asahi-machi, Kurume 830-0011, Japan. E-mail: shoichi@med.kurume-u.ac.jp

Received 29 April 2014; Accepted 25 June 2014

DOI 10.1002/jcla.21797

Published online in Wiley Online Library (wileyonlinelibrary.com).

often detected at an advanced stage, and NBNC-HCC patients were reported to have a poorer prognosis than hepatitis virus-related HCC subjects (3–5, 9, 10).

Nonalcoholic fatty liver (NAFL) is the most common chronic liver disease in the world. According to the annual health-check reports in Japan, 9–30% of Japanese adults have suffered from NAFL due to the Westernization of lifestyles and increasing rates of obesity and diabetes (11–15). NAFL is characterized by hepatic steatosis in the absence of significant alcohol intake or other known liver diseases. NAFL includes a wide spectrum of liver diseases, ranging from fatty liver, a benign and nonprogressive condition, to nonalcoholic steatohepatitis (NASH), which can progress to liver cirrhosis and HCC (16–19). Indeed, several case series of NASH-associated HCC have been actually reported (20, 21).

Advanced glycation end products (AGE) were originally characterized by a yellow-brown fluorescent color and an ability to form cross-links with and between amino groups, but the term is now used for a broad range of advanced products of the glycation process, also called the Maillard reaction (22–24). The formation and accumulation of AGE in various tissues have been known to progress during normal aging, and at an extremely accelerated rate in diabetes mellitus (22–24). Recent studies have suggested that AGE can be formed not only from reducing sugars, but also from carbonyl compounds derived from the auto-oxidation of sugars and the other metabolic pathways (25, 26). There is accumulating evidence to show that AGE play a role in the pathogenesis of a variety of disorders such as diabetic vascular complications, alcoholic liver injury, NASH, Alzheimer's disease, osteoporosis and cancer growth, and metastasis (27–34). However, the clinical significance for measuring serum AGE levels as a biomarker of NBNC-HCC remains unknown. Therefore, in this study, we examined whether serum levels of AGE were elevated in NBNC-HCC patients compared with NASH subjects without HCC or controls and further investigated which anthropometric and clinical variables were independent determinants of AGE.

PATIENTS AND RESEARCH DESIGN

Ninety patients with treatment-naïve NBNC-HCC were enrolled between April 2008 and March 2012. In addition, 27 volunteers and 56 NASH without HCC were enrolled as controls and comparators for NBNC-HCC, respectively. For the enrollment of volunteers, steatosis, liver diseases, and metabolic complications were excluded by ultrasound scan and by clinical laboratory examination. All had normal physical examinations and liver function tests were negative for serology and viral hepatitis

and had no history of liver diseases. In NASH patients, current and past daily alcohol intake was less than 20 g/week. None of the patients had received any medication that could cause NASH. Among these patients, those with the following disorders were excluded: secondary cause of steatohepatitis and drug-induced liver disease, alcohol liver disease, viral hepatitis, autoimmune hepatitis, primary biliary cirrhosis, α_1 -antitrypsin deficiency, hemochromatosis, Wilson's disease, and biliary obstruction (29). Histological diagnosis for NAFLD was performed according to the methods of Matteoni et al. (18). NBNC-HCC was defined as negative for both HBsAg and HCVAb. HBsAg in sera were assayed using the HBsAg-EIA Cobas Core Test (Hoffmann La Roche). Anti-HCV in sera was assayed using the Anti-HCV-EIA Cobas Core Test (Hoffmann La Roche, Basel, Switzerland). Patients with autoimmune hepatitis or with primary biliary cirrhosis were also excluded from NBNC-HCC group. Heavy alcohol consumption was defined when alcohol intake was more than 80 g/day for 5 years (35). Etiology of NBNC-HCC patients with heavy alcohol consumption was estimated to be alcohol. Informed consent was obtained from all patients, and the study was conducted in conformity with the ethical guidelines of the seventh revision of the Declaration of Helsinki (36) and was approved by the ethics and research committees of Hiroshima University Hospital.

The body mass index was calculated as the weight (kg) divided by height (m)-squared. Blood pressure was measured in the sitting position using an upright standard sphygmomanometer. Venous blood samples were taken in the morning following overnight fasting for 12 h. Blood chemistries such as aspartate aminotransferase (AST), alanine aminotransferase (ALT), γ -glutamyl transpeptidase (GGT), fasting plasma glucose (FPG), glycated hemoglobin (HbA1c), and lipid parameters were measured using the standard methods as described previously (29). Insulin resistance was calculated by the homeostasis model assessment of insulin resistance (HOMA-IR) using the following formula: $\text{HOMA-IR} = \text{fasting insulin } (\mu\text{U/ml}) \times \text{plasma glucose (mg/dl)} / 405$. For the measurements of AGE, serum was stored at -30°C until use. Serum levels of AGE was measured with a competitive enzyme-linked immunosorbent assay system as described previously (29). All measurements were made in triplicate, and the average values were used in the statistical analyses.

Data are presented as mean $\pm$ standard deviation (SD) or as number. All analysis was performed using the R statistical package (<http://www.r-project.org>). Statistical differences in quantitative data were determined using the Mann–Whitney *U*-test, the Kruskal–Wallis test or Chi-squared test. Differences were considered statistically significant at *P*-values less than 0.05.

TABLE 1. Characteristics of Patients

Characteristics	NBNC-HCC (<i>n</i> = 90)	NASH without HCC (<i>n</i> = 56)	Control (<i>n</i> = 27)
Sex (male/female)	71/19*	39/17	13/14
Age (year)	69.3 ± 9.7*,**	49.9 ± 12.9	48.4 ± 12.0
Body mass index (kg/m ²)	24.1 ± 3.6*,**	27.1 ± 3.7*	21.8 ± 2.2
AST (U/l)	59.1 ± 71.2*	53.4 ± 35.3*	20.5 ± 4.8
ALT (U/l)	49.5 ± 105.3**	93.9 ± 72.4*	17.1 ± 6.9
GGT (U/l)	157.7 ± 187.8*,**	88.4 ± 98.9*	28.9 ± 16.3
LDL-C (mg/dl)	103.9 ± 33.7**	142.3 ± 27.4*	109.0 ± 15.8
TG (mg/dl)	107.5 ± 60.1**	184.7 ± 86.5*	80.1 ± 37.4
HDL-C (mg/dl)	48.5 ± 14.8*	49.0 ± 12.8*	65.2 ± 17.4
FPG (mg/dl)	125.6 ± 42.3*,**	107.9 ± 18.7*	96.7 ± 7.5
HbA1c (%)	6.0 ± 1.3*	5.7 ± 0.9	5.3 ± 0.4
HOMA-IR	3.4 ± 2.5*	3.2 ± 2.0*	1.8 ± 1.2
AGE (U/ml)	9.1 ± 2.7*,**	5.2 ± 1.7*	3.5 ± 1.2
Systolic blood pressure (mmHg)	118.4 ± 8.0*	125.0 ± 14.5*	110.4 ± 13.4
Diastolic blood pressure (mmHg)	72.0 ± 7.8	74.8 ± 11.2	69.4 ± 7.2

P* < 0.05 compared with control. *P* < 0.05 compared with NASH.

RESULTS AND DISCUSSION

Clinical and laboratory characteristics of 90 patients with treatment-naïve NBNC-HCC, 56 biopsy-proven NASH and 27 healthy controls are shown in Table 1. Among NBNC-HCC patients, 10 were NASH-associated HCC, 49 alcoholic liver disease-associated one, and the etiology of rest of them (*N* = 31) was unknown. As shown in Table 1, serum levels of AGE in NBNC-HCC patients were significantly higher than those in NASH subjects without HCC or controls (9.1 ± 2.7 vs. 5.2 ± 1.7 or 3.5 ± 1.2 U/ml, respectively, *P* < 0.05). There was no significant difference in serum AGE levels among NASH-, alcohol-associated and unknown-etiology HCC (9.1 ± 2.1 vs. 8.6 ± 1.9 vs. 9.8 ± 3.7 U/ml). Compared with NASH patients without HCC, age, GGT, and FPG were significantly higher in NBNC-HCC, while body mass index, ALT, low-density lipoprotein-cholesterol (LDL-cholesterol), and triglycerides (TG) were lower. As the case of NASH subjects without HCC, compared with controls, age, GGT and FPG were significantly elevated in NBNC-HCC patients. Although there was no significant difference of ratio of male to female, AST, high-density lipoprotein-cholesterol (HDL-cholesterol), HbA1c, and HOMA-IR between NBNC-HCC and NASH subjects without HCC, ratio of male to female, AST, HbA1c, and HOMA-IR were significantly higher, whereas HDL-cholesterol was lower in NBNC-HCC patients compared with controls.

Univariate analysis revealed that male (*P* < 0.05), age (*P* < 0.01), AST (*P* < 0.05), GGT (*P* < 0.01), HDL-cholesterol (inversely, *P* < 0.01), FPG (*P* < 0.01), and HbA1c (*P* < 0.05) were significantly associated with serum levels of AGE (Table 2). Because these significant parameters could be closely correlated with each other,

we performed multiple stepwise regression analysis in order to determine the independent correlates of serum AGE levels. As shown in Table 2, age, GGT, and HDL-cholesterol (inversely) remained significant and were independently related to AGE levels ($R^2 = 0.406$).

We have previously shown that AGE could play a role in the pathogenesis of NASH and alcoholic liver injury in humans (28–30, 37). Indeed, in vitro, AGE could stimulate the proliferation and activation of hepatic stellate cells, thereby causing hepatic inflammation and fibrosis (37). In humans, serum levels of AGE were significantly elevated in NASH patients compared with simple steatosis (29). Furthermore, chronic ethanol ingestion has been shown to stimulate acetaldehyde-induced AGE formation in the liver and correlate with the severity of alcoholic liver disease in rats (30). In this study, we found for the first time that serum levels of AGE were significantly higher compared with NASH subjects without HCC or controls. The present findings have extended our previous observations showing that (1) AGE not only enhance the angiogenic potential of HCC by upregulating vascular endothelial growth factor expression, but also stimulate the proliferation of HCC in vitro (38). AGE might contribute to the development and progression of NBNC-HCC in humans, thereby being a biomarker of NBNC-HCC.

In the present study, besides age, GGT and HDL-cholesterol (inversely) were independent correlates of serum AGE levels in all subjects. GGT levels have been shown to be a biomarker that could predict outcomes and overall survival in HCC patients treated with or without transarterial chemoembolization or operation (39–42). These findings suggest that increased AGE levels could partly explain the link between high GTT values and shorter survival in HCC patients. Further, since HDL-cholesterol levels were reported to decrease in HCC

TABLE 2. Univariate and Multivariate Stepwise Regression Analysis for the Determinants of AGE

Characteristics	Univariate		Multivariate	
	β	<i>P</i>	β	<i>P</i>
Sex (male/female)	−0.161	<i>P</i> < 0.05		
Age (year)	0.550	<i>P</i> < 0.01	0.518	<i>P</i> < 0.01
Body mass index (kg/m ²)	−0.036	0.638		
AST (U/L)	0.170	<i>P</i> < 0.05		
ALT (U/L)	0.052	0.508		
GGT (U/L)	0.271	<i>P</i> < 0.01	0.206	<i>P</i> < 0.01
LDL-C (mg/dl)	−0.095	0.232		
TG (mg/dl)	−0.070	0.377		
HDL-C (mg/dl)	−0.235	<i>P</i> < 0.01	−0.253	<i>P</i> < 0.01
FPG (mg/dl)	0.257	<i>P</i> < 0.01		
HbA1c (%)	0.193	<i>P</i> < 0.05		
HOMA-IR	0.006	0.951		
Systolic blood pressure (mmHg)	0.013	0.900		
Diastolic blood pressure (mmHg)	−0.024	0.826		

Male = 0, Female = 1. $R^2 = 0.406$

subjects (43), AGE might decrease HDL-cholesterol levels, which could lead to poor prognosis in these patients.

CONCLUSIONS

In this study, we found that circulating AGE levels were significantly increased in NBNC-HCC patients compared with NASH subjects without HCC or controls. The present results suggest that AGE could be involved in the pathogenesis of NBNC-HCC, thereby being a biomarker that might discriminate NBNC-HCC from NASH.

ACKNOWLEDGMENTS

This study was supported in part by Grants-in-Aid for Scientific Research (B) from the Ministry of Education, Culture, Sports, Science and Technology, Japan (to M.T. and to S.Y.), and by MEXT-Supported Program for the Strategic Research Foundation at Private Universities, the Ministry of Education, Culture, Sports, Science and Technology (MEXT) (to S.Y.).

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

- Bosch FX, Ribes J, Díaz M, Cléries R. Primary liver cancer: World-wide incidence and trends. *Gastroenterology* 2004;127(5 Suppl 1):S5–S16.
- Siegel R, Naishadham D, Jemal A. Cancer statistics, 2012. *CA Cancer J Clin* 2012;62:10–29.
- Takuma Y, Nouse K, Makino Y, et al. Outcomes after curative treatment for cryptogenic cirrhosis-associated hepatocellular carcinoma satisfying the Milan criteria. *J Gastroenterol Hepatol* 2011;26:1417–1424.
- Kaibori M, Ishizaki M, Matsui K, Kwon AH. Clinicopathologic characteristics of patients with non-B non-C hepatitis virus hepatocellular carcinoma after hepatectomy. *Am J Surg* 2012;204:300–307.
- Akahoshi H, Taura N, Ichikawa T, et al. Differences in prognostic factors according to viral status in patients with hepatocellular carcinoma. *Oncol Rep* 2010;23:1317–1323.
- Nagaoki Y, Hyogo H, Aikata H, et al. Recent trend of clinical features in patients with hepatocellular carcinoma. *Hepatol Res* 2012;42:368–375.
- Umemura T, Kiyosawa K. Epidemiology of hepatocellular carcinoma in Japan. *Hepatol Res* 2007;37 Suppl 2:S95–S100.
- Hatanaka K, Kudo M, Fukunaga T, et al. Clinical characteristics of NonBNonC- HCC: Comparison with HBV and HCV related HCC. *Intervirology* 2007;50:24–31.
- El-Serag HB. Epidemiology of viral hepatitis and hepatocellular carcinoma. *Gastroenterology* 2012;142:1264–1273.
- Yang JD, Harmsen WS, Slettedahl SW, et al. Factors that affect risk for hepatocellular carcinoma and effects of surveillance. *Clin Gastroenterol Hepatol* 2011;9: 617–623.
- Hamaguchi M, Kojima T, Takeda N, et al. The metabolic syndrome as a predictor of nonalcoholic fatty liver disease. *Ann Intern Med* 2005;143:722–728.
- Kojima S, Watanabe N, Numata M, Ogawa T, Matsuzaki S. Increase in the prevalence of fatty liver in Japan over the past 12 years: Analysis of clinical background. *J Gastroenterol* 2003;38: 954–961.
- Amarapurkar DN, Hashimoto E, Lesmana LA, Sollano JD, Chen PJ, Goh KL; Asia-Pacific Working Party on NAFLD. How common is non-alcoholic fatty liver disease in the Asia-Pacific region and are there local differences? *J Gastroenterol Hepatol* 2007;22:788–793.
- Okanoue T, Umemura A, Yasui K, Itoh Y. Nonalcoholic fatty liver disease and nonalcoholic steatohepatitis in Japan. *J Gastroenterol Hepatol* 2011;26 Suppl 1:153–162.
- Eguchi Y, Hyogo H, Ono M, et al. JSG-NAFLD. Prevalence and associated metabolic factors of nonalcoholic fatty liver disease in the general population from 2009 to 2010 in Japan: A multicenter large retrospective study. *J Gastroenterol* 2012;47:586–595.

16. Teli MR, James OF, Burt AD, Bennett MK, Day CP. The natural history of nonalcoholic fatty liver: a follow-up study. *Hepatology* 1995;22:1714–1719.
17. Dam-Larsen S, Franzmann M, Andersen IB, et al. Long term prognosis of fatty liver: risk of chronic liver disease and death. *Gut* 2004;53:750–755.
18. Matteoni CA, Younossi ZM, Gramlich T, Boparai N, Liu YC, McCullough AJ. Nonalcoholic fatty liver disease: a spectrum of clinical and pathological severity. *Gastroenterology* 1999;116:1413–1419.
19. James O, Day C. Non-alcoholic steatohepatitis: another disease of affluence. *Lancet* 1999;353:1634–1636.
20. Starley BQ, Calcagno CJ, Harrison S A., Nonalcoholic fatty liver disease and hepatocellular carcinoma: A weighty connection. *Hepatology* 2010;51:1820–1832.
21. Takuma Y, Noso K. Nonalcoholic steatohepatitis-associated hepatocellular carcinoma: our case series and literature review. *World J Gastroenterol* 2010;16:1436–1441.
22. Brownlee M, Cerami A, Vlassara H. Advanced glycosylation end products in tissue and the biochemical basis of diabetic complications. *N Engl J Med* 1988;318:1315–1321.
23. Al-Abed Y, Kapurniotu A, Bucala R. Advanced glycation end products: Detection and reversal. *Methods Enzymol* 1999;309:152–172.
24. Vlassara H, Palace MR. Diabetes and advanced glycation endproducts. *J Intern Med* 2002;251:87–101.
25. Glomb MA, Monnier VM. Mechanism of protein modification by glyoxal and glycolaldehyde, reactive intermediates of the Maillard reaction. *J Biol Chem* 1995;270:10017–10026.
26. Thornalley PJ, Langborg A, Minhas HS. Formation of glyoxal, methylglyoxal and 3-deoxyglucosone in the glycation of proteins by glucose. *Biochem J* 1999;344:109–116.
27. Yamagishi S, Maeda S, Matsui T, Ueda, S, Fukami K, Okuda S. Role of advanced glycation end products (AGEs) and oxidative stress in vascular complications in diabetes. *Biochim Biophys Acta* 2012;1820:663–671.
28. Hyogo H, Yamagishi S, Advanced glycation end products (AGEs) and their involvement in liver disease. *Curr Pharm Des* 2008, 14, 969–972.
29. Hyogo H, Yamagishi S, Iwamoto K, et al. Elevated levels of serum advanced glycation end products in patients with non-alcoholic steatohepatitis. *J Gastroenterol Hepatol* 2007;22:1112–1119.
30. Hayashi N, George J, Takeuchi M, et al. Acetaldehyde-derived advanced glycation end-products promote alcoholic liver disease. *PLoS One* 2013;8:e70034.
31. Takino J, Yamagishi S, Takeuchi M. Glycer-AGEs-RAGE signaling enhances the angiogenic potential of hepatocellular carcinoma by upregulating VEGF expression. *World J Gastroenterol* 2012;18:1781–1788.
32. Yamagishi S. Role of advanced glycation end products (AGEs) in osteoporosis in diabetes. *Curr Drug Targets* 2011;12:2096–2102.
33. Takeuchi M, Yamagishi S. Possible involvement of advanced glycation end-products (AGEs) in the pathogenesis of Alzheimer's disease. *Curr Pharm Des* 2008;14:973–978.
34. Abe R, Yamagishi S. AGE-RAGE system and carcinogenesis. *Curr Pharm Des* 2008;14:940–945.
35. Takeda A, A Japanese study group for alcoholic liver disease. A new diagnostic criteria of alcoholic liver disease. *Kanzo* 1993;34:888–896.
36. The world medical association, inc. Declaration of Helsinki; Ethical principle for medical research involving human subject. WMA General Assembly, Seoul, Korea, October 2008.
37. Iwamoto K, Kanno K, Hyogo H, et al. Advanced glycation end products enhance the proliferation and activation of hepatic stellate cells. *J Gastroenterol* 2008;43:298–304.
38. Sakurao Y, Sawada T, Okada T, et al. MK615 decreases RAGE expression and inhibits TAGE-induced proliferation in hepatocellular carcinoma cells. *World J Gastroenterol* 2010;16:5334–5341.
39. Guiu B, Deschamps F, Boulin M, et al. Serum gamma-glutamyl-transferase independently predicts outcome after transarterial chemoembolization of hepatocellular carcinoma: External validation. *Cardiovasc Intervent Radiol* 2012;35:1102–1108.
40. Zhang JB, Chen Y, Zhang B, et al. Prognostic significance of serum gamma-glutamyl transferase in patients with intermediate hepatocellular carcinoma treated with transcatheter arterial chemoembolization. *Eur J Gastroenterol Hepatol* 2011;23:787–793.
41. Ju MJ, Qiu SJ, Fan J, et al. Preoperative serum gamma-glutamyl transferase to alanine aminotransferase ratio is a convenient prognostic marker for Child-Pugh A hepatocellular carcinoma after operation. *J Gastroenterol* 2009;44:635–642.
42. Morsi MI, Hussein AE, Mostafa M, El-Abd E, El-Moneim NA. Evaluation of tumour necrosis factor-alpha, soluble P-selectin, gamma-glutamyl transferase, glutathione S-transferase-pi and alpha-fetoprotein in patients with hepatocellular carcinoma before and during chemotherapy. *Br J Biomed Sci* 2006;63:74–78.
43. Jiang J, Nilsson-Ehle P, Xu N. Influence of liver cancer on lipid and lipoprotein metabolism. *Lipids Health Dis* 2006;5:4.